

Leptogenesis from spin-gravity coupling following inflation

Subhendra Mohanty^a, A.R.Prasanna^a, and G. Lambiase^{b,c}

^aPhysical Research Laboratory, Navrangpura, Ahmedabad - 380 009, India.

^bDipartimento di Fisica "E.R. Caianiello" Università di Salerno, 84081 Baronissi (Sa), Italy. and

^cINFN - Gruppo Collegato di Salerno, Italy.

The energy levels of the left and the right handed neutrinos is split in the background of gravitational waves generated during inflation which, in presence of lepton number violating interactions, gives rise to a net lepton asymmetry at equilibrium. Lepton number violation is achieved by the same dimension five operator which gives rise to neutrino masses after electro-weak symmetry breaking. A net baryon asymmetry of the same magnitude can be generated from this lepton asymmetry by electroweak sphaleron processes.

PACS numbers: 04.30.Db, 04.62.+v, 98.80.Cq

The successful prediction of light element abundance by the big-bang nucleosynthesis [1] depends on the assumption that the net baryon number to entropy ratio η as determined by the recent WMAP [2] results is $\eta = 9.2^{+0.6}_{-0.4} \times 10^{-11}$. Sakharov's well known condition demand that in a *CPT* conserving theory, a net lepton/baryon asymmetry can be generated if there is (a) *B/L* violating interactions (b) *C* and *CP* violation so that *B/L* violating reactions in the forward and reverse channels do not cancel and (c) departure from thermal equilibrium as the statistical distribution of particles and anti-particles is the same if the Hamiltonian commutes with *CPT*. If in the theory we have *CPT* is violation it is possible to generate lepton/baryon asymmetry at thermal equilibrium without requiring *CP* violation.

In this paper we show that *CPT* is violated spontaneously due to the spin-connection couplings of fermions with cosmological gravitational waves. It is well known that inflation [3] generates a nearly scale invariant spectrum of gravitational waves [4]. The spin connection couplings split the energy levels of neutrinos compared to anti-neutrinos and in presence of lepton number violating interaction there is a net asymmetry generated between neutrinos and anti-neutrinos at *thermodynamic equilibrium*. Lepton number violation is generated by the dimension five operator introduced by Weinberg [5] which also generates the neutrino masses after electro-weak symmetry breaking. The lepton-asymmetry gets frozen-in when the lepton-number violating processes decouple. Baryon asymmetry can then be generated from this lepton-asymmetry by the electro-weak sphaleron processes [6]. Sphaleron processes conserve (*B - L*) so a lepton asymmetry generated in the GUT era can be converted to baryon asymmetry of the same magnitude [7].

The general covariant coupling of spin 1/2 particles to gravity is given by the Lagrangian [8] $\mathcal{L} = \sqrt{-g}(\bar{\psi}\gamma^a D_a \psi - m\bar{\psi}\psi)$, $D_a = \partial_a - \frac{i}{4}\omega_{bca}\sigma^{bc}$ is the covariant derivative, and ω_{bca} are the spin-connections $\omega_{bca} = e_{b\lambda}(\partial_a e^\lambda_c + \Gamma^\lambda_{\gamma\mu} e^\gamma_c e^\mu_a)$. This Lagrangian is invariant under the local Lorentz transformation of the vierbein $e^\mu_a(x) \rightarrow \Lambda^\mu_b(x)e^\mu_b(x)$ and the spinor fields $\psi(x) \rightarrow \exp(i\epsilon_{ab}(x)\sigma^{ab})\psi(x)$. Here $\sigma^{bc} = \frac{i}{2}[\gamma^b, \gamma^c]$ are genera-

tors of tangent space Lorentz transformation (*a, b, c* etc. denote the inertial frame 'flat space' indices and α, β, γ etc. are the coordinate frame 'curved space' indices such that $e^\mu_a e^{\nu a} = g^{\mu\nu}$, $e^{a\mu} e^\mu_b = \eta^{ab}$, $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$ where η^{ab} represents the inertial frame Minkowski metric, and $g_{\mu\nu}$ is the curved space metric).

The spin-connection term in the Dirac equation is a product of three Dirac matrices which after some algebra can be reduced to a vector $A^a \gamma_a$ and an axial vector $iB^d \gamma_5 \gamma_d$. The vector term turns out to be anti-hermitian and disappears when the hermitian conjugate part is added to the lagrangian $\mathcal{L}$. The surviving interaction term which describes the spin-connection coupling of fermions to gravity can be written as an axial-vector

$$\begin{aligned} \mathcal{L} &= \det(e) \bar{\psi} (i\gamma^a \partial_a - m - \gamma_5 \gamma_d B^d) \psi, \\ B^d &= \epsilon^{abcd} e_{b\lambda} (\partial_a e^\lambda_c + \Gamma^\lambda_{\alpha\mu} e^\alpha_c e^\mu_a). \end{aligned} \quad (1)$$

In a local inertial frame of the fermion, the effect of a gravitational field appears as an axial-vector interaction term shown in (1). We now calculate the four vector B^d for a perturbed Robertson-Walker universe.

The general form of perturbations on a flat Robertson-Walker expanding universe can be written as [9]

$$\begin{aligned} ds^2 &= a(\tau)^2 [(1 + 2\phi)d\tau^2 - \omega_i dx^i d\tau \\ &\quad - ((1 + 2\psi)\delta_{ij} + h_{ij})dx^i dx^j], \end{aligned} \quad (2)$$

where ϕ and ψ are scalar, ω_i are vector and h_{ij} are the tensor fluctuations of the metric. Of the ten degrees of freedom in the metric perturbations only six are independent and the remaining four can be set to zero by suitable gauge choice. For our application we need only the tensor perturbations and we choose the transverse-traceless (TT) gauge $h^i_i = 0, \partial^i h_{ij} = 0$ for the tensor perturbations. In the TT gauge the perturbed Robertson-Walker metric can be expressed as

$$\begin{aligned} ds^2 &= a(\tau)^2 [(1 + 2\phi)d\tau^2 - \omega_i dx^i d\tau - (1 + 2\psi - h_+)dx_1^2 \\ &\quad - (1 + 2\psi + h_+)dx_2^2 - 2h_\times dx_1 dx_2 - (1 + 2\psi)dx_3^2]. \end{aligned} \quad (3)$$

An orthogonal set of vierbiens e_μ^a for this metric is given by

$$e_\mu^a = a(\tau) \begin{pmatrix} 1 + \phi & -\omega_1 & -\omega_2 & -\omega_3 \\ 0 & -(1 + \psi) + h_+/2 & h_\times & 0 \\ 0 & 0 & -(1 + \psi) - h_+/2 & 0 \\ 0 & 0 & 0 & -(1 + \psi) \end{pmatrix}. \quad (4)$$

Using the vierbiens (4) the expression for the components of the four vector field B^d (1) is given by

$$B^0 = \partial_3 h_\times, \quad B^i = (\nabla \times \vec{\omega})^i + \partial_\tau h_\times \delta^{i3}. \quad (5)$$

The choice of vierbiens (4) which gives the metric (2) is not unique as one can make a local Lorentz transformation (LLT) $e_\mu^{m'} = \Lambda_n^{m'} e_\mu^n$. Under a LLT the four-vector B^d transforms as $B^{d'} = \Lambda_a^{d'} B^a$. The dispersion relations of the left and the right helicity fermions have B^d dependent terms of the form $\eta_{mn} B^m B^n$ and $\eta_{mn} p^m B^n$, and therefore the dispersion relations do not change with a transformation of the local inertial frame.

The fermion bilinear term $\bar{\psi} \gamma_5 \gamma_a \psi$ is odd under CPT transformation. When one treats B^a as a background field then the interaction term in (1) explicitly violates CPT . When the primordial metric fluctuations become classical, i.e there is no back-reaction of the micro-physics involving the fermions on the metric and B^a is considered as a fixed external field, then CPT is violated spontaneously.

The gravitational spin connection coupling for the neutrinos at high energy is given by

$$\mathcal{L} = \det(e) [(i\bar{\nu}_L \gamma^a \partial_a \nu_L + i\bar{\nu}_R \gamma_a \partial_a \nu_R) + m\bar{\nu}_L \nu_R + m^\dagger \bar{\nu}_R \nu_L + B^a (\bar{\nu}_R \gamma_a \nu_R - \bar{\nu}_L \gamma_a \nu_L)], \quad (6)$$

where B^a are the parameters of the gravitational waves as defined in (5). If we consider only the Standard Model fermions then the right handed neutrinos carry the opposite Lepton number compared to the left handed neutrinos, $\nu_R = (\nu_L)^c$ and the mass term in (6) is of the Majorana type (we have suppressed the generation index).

The dispersion relation of left and right helicity neutrinos fields are given by $\eta^{ab}(p_a + \xi B_a)(p_b + \xi B_b) = m^2$, where $\xi = -1$ for ν_L and $\xi = 1$ for ν_R . Keeping terms linear in the perturbations B^a , the free particle energy of the left and right helicity states is

$$E_{L,R}(p) = p + \frac{m^2}{2p} \mp \left(B_0 - \frac{\mathbf{p} \cdot \mathbf{B}}{p} \right), \quad (7)$$

with $p = |\mathbf{p}|$. In the Standard Model ν_L carry lepton number +1 and ν_R are assigned lepton number (-1). In the presence of non-zero metric fluctuations, there is a split in energy levels of $\nu_{L,R}$ given by (7). If there are GUT processes that violate lepton number freely above

some decoupling temperature T_d , then the equilibrium value of lepton asymmetry generated for all $T > T_d$ will be

$$n(\nu_L) - n(\nu_R) = \frac{g}{2\pi^2} \int d^3p \left[\frac{1}{1 + e^{\frac{E_L}{T}}} - \frac{1}{1 + e^{\frac{E_R}{T}}} \right] \quad (8)$$

The spin-connection coupling with gravitational waves also splits the energy levels between the charged left and right handed fermions. For example $E(e_R^-) - E(e_L^-) = 2(B_0 - \mathbf{p} \cdot \mathbf{B}/p)$. But this does not lead to lepton generation of lepton asymmetry as both e_L^- and e_R^- carry the same lepton number.

In the ultra-relativistic regime $p \gg m_\nu$ and assuming that $B_0 \ll T$, the expression (8) for lepton asymmetry reduces to

$$\Delta n_L = \frac{gT^3}{6} \left(\frac{B_0}{T} \right). \quad (9)$$

The dependence on $\mathbf{B}$ drops out after angular integration in (8) and the lepton asymmetry depends on the tensor perturbations only through B^0 .

To compute the spectrum of gravitational waves $h(\mathbf{x}, \tau)$ during inflation, we express $h_\times$ in terms of the creation- annihilation operator

$$h(\mathbf{x}, \tau) = \frac{\sqrt{16\pi}}{aM_p} \int \frac{d^3k}{(2\pi)^{3/2}} \left(a_{\mathbf{k}} f_k(\tau) + a_{-\mathbf{k}}^\dagger f_k^*(\tau) \right) e^{i\mathbf{k} \cdot \mathbf{x}} \quad (10)$$

where $\mathbf{k}$ is the comoving wavenumber, $k = |\mathbf{k}|$, and $M_p = 1.22 \cdot 10^{19} \text{ GeV}$ is the Planck mass. The mode functions $f_k(\tau)$ obey the minimally coupled Klein-Gordon equation

$$f_k'' + \left(k^2 - \frac{a''}{a} \right) f_k = 0. \quad (11)$$

During de Sitter era, the scale factor $a(\tau) = -1/(H_I \tau)$ where H_I is the Hubble parameter, and Eq. (11) has the solution

$$f_k(\tau) = \frac{e^{-ik\tau}}{\sqrt{2k}} \left(1 - \frac{i}{k\tau} \right), \quad (12)$$

which matches the positive frequency "flat space" solutions $e^{-ik\tau}/\sqrt{2k}$ in the limit of $k\tau \gg 1$. The first term of (12) represents the decaying part of h and can be dropped. The second term of (12) represents the amplitude constant gravitational wave, which survives to

the present era. Substituting the second term of (12) in (10) and using the canonical commutation relation for a_k and $a_k^\dagger$ we get the standard expression for two point correlation of gravitational waves generated by inflation

$$\langle h(\mathbf{x}, \tau) h(\mathbf{x}, \tau) \rangle^{inf} \equiv \int \frac{dk}{k} (|h_k|^2)^{inf}, \quad (13)$$

with the spectrum of gravitational waves given by the scale invariant form

$$(|h_k|^2)^{inf} = \frac{4}{\pi} \frac{H_I^2}{M_p^2}. \quad (14)$$

There is a stringent constraint $H_I/M_p < 10^{-5}$ from CMB data [10]. This constraint limits the parameter space of interactions that can be used for generating the requisite lepton-asymmetry. In the radiation era, when these modes re-enter the horizon, the amplitude redshifts by a^{-1} from the time of re-entry. The reason is that in the radiation era $a(\tau) \sim \tau$ and the equation for f_k , (11), gives plane wave solutions $f_k = (1/\sqrt{2k}) \exp(-ik\tau)$. Therefore in the radiation era the amplitudes of h redshifts as a^{-1} . The gravitational waves inside the horizon in the radiation era will be

$$h_k^{rad} = h_k^{inf} \frac{a_k}{a(\tau)} = h_k^{inf} \frac{T}{T_k}, \quad (15)$$

where h_k^{inf} are the gravitational waves generated by inflation (14), a_k and T_k are the scale factor and the temperature when the modes of wavenumber k entered the horizon in the radiation era. The horizon entry of mode k occurs when

$$\frac{a_k H_k}{k} = \frac{a(T) T H_k}{T_k k} = 1, \quad (16)$$

where $H_k = 1.67 \sqrt{g_*} T_k^2 / M_p$ is the Hubble parameter at the time of horizon crossing of the k the mode (g_* is the number of relativistic degrees of freedom which for the Standard Model is $g_* = 106.7$). Solving equation (16) for T_k we get

$$T_k = \frac{1}{1.67 \sqrt{g_*}} \frac{k M_p}{a(\tau) T}. \quad (17)$$

The amplitude of the gravitational waves of mode k inside the radiation horizon is, using (17) and (15), given by $h_k^{rad} = h_k^{inf} \frac{a(T)}{k} \frac{T^2 1.67 \sqrt{g_*}}{M_p}$. Note that the gravitational wave spectrum inside the radiation era horizon is no longer scale invariant. The gravitational waves in position space have the correlation function

$$\langle h(\mathbf{x}, \tau) h(\mathbf{x}, \tau) \rangle^{rad} = \int \frac{dk}{k} (h_k^{rad})^2, \quad (18)$$

and hence for the spin connection B^0 generated by the inflationary gravitational waves in the radiation era, we

get

$$\begin{aligned} \langle B^0(\mathbf{x}, \tau) B^0(\mathbf{x}, \tau) \rangle &= \int \frac{dk}{k} \left(\frac{k}{a} \right)^2 (h_k^{rad})^2 \\ &= \frac{4}{\pi} \left(\frac{H_I}{M_p^2} T^2 1.67 \sqrt{g_*} \right)^2 \int_{k_{min}}^{k_{max}} \frac{dk}{k} \end{aligned} \quad (19)$$

Now we find that the spectrum of spin-connection is scale invariant inside the radiation horizon. This is significant in that the lepton asymmetry generated by this mechanism depends upon the infrared and ultraviolet scales only logarithmically. The scales outside the horizon are blue-tilted which means that there will be a scale dependent anisotropy in the lepton number correlation at two different space-time points $\langle \Delta L(r) \Delta L(r') \rangle \sim A k^n$, $n > 0$, where $\Delta L(r) \equiv L(r) - \bar{L}$, is the anisotropic deviation from the mean value. Unlike in the case of CMB, this anisotropy in the lepton number is unlikely to be accessible to experiments. Nucleosynthesis calculations only give us an average value at the time of nucleosynthesis (when $T \sim 1 \text{ MeV}$). The maximum value of k are for those modes which leave the de Sitter horizon at the end of inflation. If inflation is followed by radiation domination era starting with the re-heat temperature T_{RH} then the maximum value of k in the radiation era (at temperature T) is given by $k_{max}/a(T) = H_I \left(\frac{T}{T_{RH}} \right)$. The lower limit of k is $k_{min} = e^{-N} k_{max}$ which are the modes which left the de-Sitter horizon in the beginning of inflation (N is the total e-folding of the scale factor during inflation, $N \simeq 55 - 70$). The integration over k then yields just the factor $\ln(k_{max}/k_{min}) = N$. The r.m.s value of spin connection that determines the lepton asymmetry through equation (9) is $(B_0)_{rms} = \sqrt{\langle B_0^2 \rangle}$,

$$(B_0)_{rms} = \frac{2}{\sqrt{\pi}} \left(\frac{H_I}{M_p^2} T^2 1.67 \sqrt{g_*} \right) \sqrt{N}. \quad (20)$$

The lepton asymmetry (9) as a function of temperature can therefore be expressed as (taking $g = 3$ for the three neutrino flavors)

$$\Delta n_L(T) = \frac{1}{\sqrt{\pi}} (1.67 \sqrt{g_*}) \sqrt{N} \left(\frac{T^4 H_I}{M_p^2} \right). \quad (21)$$

The lepton number to entropy density ($s = 0.44 g_* T^3$) is given by

$$\Delta L \equiv \frac{\Delta n_L(T)}{s(T)} \simeq 2.14 \frac{T H_I \sqrt{N}}{M_p^2 \sqrt{g_*}}. \quad (22)$$

Lepton number asymmetry will be generated as long as the lepton number violating interactions are in thermal equilibrium. Once these reactions decouple at some decoupling temperature T_d , which we shall determine, the $\Delta n_L(T)/s(T)$ ratio remains fixed for all $T < T_d$.

To calculate the decoupling temperature of the lepton number violating processes we turn to a specific effective dimension five operator which gives rise to Majorana masses for the neutrinos introduced by Weinberg [5] $\mathcal{L}_W = \frac{C_{\alpha\beta}}{2M} (\bar{l}_{L\alpha}^c \tilde{\phi}^*) (\tilde{\phi}^\dagger l_{L\beta}) + h.c.$ where $l_{L\alpha} = (\nu_\alpha, e_\alpha^-)^T$ is the left-handed lepton doublet (α denotes the generation), $\phi = (\phi^+, \phi^0)^T$ is the Higgs doublet and $\tilde{\phi} \equiv i\sigma_2 \phi^* = (-\phi^{0*}, \phi^{-})^T$. M is some large mass scale and $C_{\alpha\beta}$ are of order unity.

The $\Delta L = 2$ interactions that result from the operator $\mathcal{L}_W$ are

$$\nu_L + \phi^0 \leftrightarrow \nu_R + \phi^0, \quad \nu_R + \phi^{0*} \leftrightarrow \nu_L + \phi^{0*}. \quad (23)$$

In the absence of the gravitational waves the forward reactions would equal the backward reactions and no net lepton number would be generated. In the presence of a background gravitational waves the energy levels of the left and right helicity neutrinos are no longer degenerate and this leads to a difference in the number density of left and right handed neutrinos of the magnitude given by equation (9) at thermal equilibrium. This process continues till the interactions (23) decouple. The decoupling temperature is estimated as follows. The cross section for the interaction $\nu_{L\alpha} + \phi^0 \leftrightarrow \nu_{R\beta} + \phi^0$ is $\sigma = \frac{|C_{\alpha\beta}|^2}{M^2} \frac{1}{\pi}$, and interaction rate $\Gamma = \langle n_\phi \sigma \rangle$ of the $\Delta L = 2$ interactions is $\Gamma = \frac{0.122}{\pi} \frac{|C_{\alpha\beta}|^2 T^3}{M^2}$. In the electroweak era, when the Higgs field in $\mathcal{L}_W$ acquires a vev , $\langle \phi \rangle = (0, v)^T$ (where $v = 174 \text{ GeV}$), this operator gives rise to a neutrino mass matrix $m_{\alpha\beta} = \frac{v^2 C_{\alpha\beta}}{M}$. We can therefore substitute the couplings $\frac{C_{\alpha\beta}}{M}$ in terms of the light left handed Majorana neutrino mass. At the decoupling temperature the interaction rate $\Gamma(T)$ falls below the expansion rate $H(T) = 1.7\sqrt{g_*} T^2/M_p$. The decoupling temperature is obtained from equation $\Gamma(T_d) = H(T_d)$ and turns out to be

$$T_d = 13.68\pi \sqrt{g_*} \frac{v^4}{m_\nu^2 M_p}, \quad (24)$$

where m_ν is the mass of the heaviest neutrino. A lower bound on the mass of the heaviest neutrino is given by atmospheric neutrino experiments [11] $m_\nu^2 > \Delta_{atm} = 2.5 \cdot 10^{-3} eV^2$ which means that the decoupling temperature has an upper bound given by $T_d = 1.3 \cdot 10^{13} (\Delta_{atm}/m_\nu^2) \text{ GeV}$. Substituting the expression (24) for T in (22), we finally obtain the formula for lepton number

$$\begin{aligned} L &= 92.0 \left(\frac{v^4 H_I}{m_\nu^2 M_p^3} \right) \sqrt{N} \\ &= 7.4 \cdot 10^{-11} \frac{H_I}{4 \cdot 10^{14} \text{ GeV}} \frac{2.5 \cdot 10^{-3} eV^2}{m_\nu^2} \frac{\sqrt{N}}{10}. \end{aligned} \quad (25)$$

As first pointed out in [7], electroweak sphalerons at the temperatures $T \sim 10^3 \text{ GeV}$ violate $B + L$ maximally and

conserve $B - L$. Therefore a lepton asymmetry generated at an earlier epoch gets converted to baryon asymmetry of the same magnitude by the electroweak sphalerons.

The input parameters needed for generating the correct magnitude of baryogenesis ($\eta \sim 10^{-10}$) are the amplitude of the $h_\times \sim 10^{-6}$ (or equivalently the curvature during inflation $H_I \sim 10^{14} \text{ GeV}$ or the scale of inflation is the GUT scale, $V^{1/4} \sim 10^{16} \text{ GeV}$ which is allowed by CMB [12, 13], neutrino Majorana mass in the atmospheric neutrino scale $m_\nu^2 \sim 10^{-3} eV^2$ [11] and duration of inflation $H_I t = N \sim 100$ needed to solve the horizon and entropy problems in the standard inflation paradigm. All these parameters are well within experimentally acceptable limits.

To summarize the mechanism of baryogenesis we propose arises in the standard Einstein's gravity where spontaneous CPT violation is caused by gravitational waves, and the $h_\times$ gravitational wave modes which give non-zero spin connection are produced in *generic* inflation scenarios (in contrast to models [14] where baryon asymmetry is created through a gravitational Chern-Simons term, which can be generated if specific CP violating terms are introduced in the inflaton potential which can give rise to birefringent circularly polarized gravitational waves).

-
- [1] S. Burles, K.M. Nollet, M.S. Turner, Phys. Rev. D **63**, 063512 (2001).
 - [2] C. Bennet et al., Ap. J. S. **148**, 1 (2003); D.Spergel et al., Ap. J. S. **148**, 175 (2003).
 - [3] A.H. Guth, Phys. Rev. D **23**, 347 (1981).
 - [4] V.F. Mukhanov, H.A. Feldman, R.H. Brandenberger, Phys. Rept. **215**, 203 (1992).
 - [5] S. Weinberg, Phys. Rev. Lett. **43** 1566 (1979).
 - [6] V.A. Kuzmin, V.A. Rubakov, M.E. Shapashnikov, Phys. Lett. **B 155**, 36 (1985).
 - [7] M. Fukugita, T. Yanagida, Phys. Lett. **174**, 45 (1992).
 - [8] J. Schwinger, *Particles, Sources and Fields*, **Vol. 1**, p 401, Addison-Wesley, (1970).
 - [9] A. R. Liddle, D. H. Lyth, *Inflation and Large-Scale Structure*, Cambridge Univ. Press, (2000).
 - [10] L.M. Krauss, M.J. White, Phys. Rev. Lett. **69**, 869 (1992); A. Melchiorri, C. J. Odman, Phys. Rev. D **67**, 021501 (2003).
 - [11] Y. Ashie et al., Phys. Rev. D **71**, 112005 (2005).
 - [12] L. Pilo, A. Riotto, A. Zaffaroni, Phys. Rev. Lett. **92**, 201303 (2004).
 - [13] L. Knox, Y. S. Song, Phys. Rev. Lett. **89**, 011303 (2002).
 - [14] S. H. S. Alexander, M. E. Peskin, M. M. Sheikh-Jabbari, arXiv:hep-th/0403069; D. H. Lyth, C. Quimbay, Y. Rodriguez, JHEP **0503**, 016 (2005).